INFLUENCE OF TRANSLATIONAL AND ROTATIONAL ENERGY ON THE RATE OF ION-MOLECULE REACTIONS NEAR 0 K: D_2^++NH_3 AND D_2^++ND_3

R. HAHN, V. ZHELYAZKOVA, D. SCHLANDER, F. MERKT, *Laboratory of Physical Chemistry, ETH Zürich, Vladimir-Prelog-Weg 1-5/10, 8093 Zürich, Switzerland*

Ion-molecule reactions are important reactions in atmospheric chemistry, astrophysics and plasma physics[1]. Reliable values for the reaction rates are key to modelling the relevant reaction networks, but are difficult to study in the collision energy range relevant for astrophysics because ions are very easily heated up by stray electric or magnetic fields. To study the reactions D_2^++NH_3 as well as D_2^++ND_3 at low collision energy, we replace D_2^+ by D_2 in a Rydberg state (of high principal quantum number $n \geq 30$). The distant Rydberg electron shields the ion core from external fields without influencing the ion-molecule reaction within its orbit. We use a Rydberg-Stark deflector to merge a beam of D_2 Rydberg molecules with a supersonic beam of NH_3 or ND_3 and tune the collision energy E_{coll} from $\approx k_{\text{B}}\cdot$ 150 mK to $\approx k_{\text{B}}\cdot$ 70 K[2]. Short ($\approx$ 20 μs) gas pulses ensure a high collision-energy resolution. We observe two reaction channels for the reaction with NH_3: NH_3^++D_2 and NH_2D^++D, but these two channels are not resolvable by mass spectrometry for D_2^++ND_3. The rate coefficients deviate strongly from Langevin rates and increase sharply below 2 K, which is explained by rotationally adiabatic capture calculations as arising from the linear Stark effect of the two inversion tunneling components of ammonia in the electric field of the ion [3,4,5]. We study the influence of the rotational temperature (14 K or 39 K) of NH_3 and ND_3 using pure ammonia beams and beams of ammonia seeded in helium and measure the rotational states populations with (2+1) REMPI spectroscopy. We compare these results with those obtained for the He^++NH_3[6] and Rg^++NH_3 (Rg=Ar, Kr, Xe) reactions[7].

[1] doi:10.1038/s42254-020-0198-0, B. A. McGuire, O. Asvany, S. Brünken, S. Schlemmer, *Nat. Rev. Phys.*, **2**, 402-410, (2020).

[2] doi:10.1103/PhysRevA.90.043403, P. Allmendinger, J. Deiglmayr, J. A. Agner, H. Schmutz, F. Merkt, *Phys. Rev. A*, **90**, 043403, (2014).

[3] doi:10.1063/1.453701, J. Troe, *Chem. Phys.*, **87**, 2773-2780, (1987).

[4] doi:10.1080/00268978500100461, D. C. Clary, *Mol. Phys.*, **54**, 605-618, (1985).

[5] doi:10.1063/1.472479, J. Troe, *J. Chem. Phys.*, **105**, 6249-6262, (1996).

[6] doi:10.1039/D1CP03116C, V. Zhelyazkova, F. B. V. Martins, J. A. Agner, H. Schmutz, F. Merkt, *Phys. Chem. Chem. Phys.*, **23**, 21606, (2021).

[7] doi:10.1039/D1SC01652K, A. Tsikritea, K. Park, P. Bertier, J. Loreau, T. Softley, B. R. Heazlewood, *Chem. Sci.*, **12**, 10005-10013, (2021).