

**ANALYSIS OF THE HOT BANDS OF METHYLAMINE: $3\nu_{15} \leftarrow \nu_{15}$,
 $4\nu_{15} \leftarrow \nu_{15}$ AND $\nu_9 \leftarrow \nu_{15}$**

I. GULACZYK, M. KREGLEWSKI, *Faculty of Chemistry, Adam Mickiewicz University in Poznan, Poland*; **I. KLEINER**, *Universite Paris Cite and Univ Paris Est Creteil, CNRS, LISA, 75013, Paris, France*

Methylamine is a molecule performing two large amplitude motions: CH_3 internal rotation and NH_2 inversion. These two large amplitude motions are strongly coupled and give rise to a rotation-inversion-torsion structure in the vibrational states. The rovibrational spectrum of the methylamine molecule has been extensively studied both experimentally and theoretically. The analyses of infrared bands such as inversion (ν_9) or CN stretching (ν_8) show significant perturbations from highly excited torsional states. In order to untangle the interactions in the 700 - 1200 cm^{-1} region of the methylamine spectrum, it is crucial to assign the dark states, which are the third and fourth excited torsional states ($3\nu_{15}$ and $4\nu_{15}$). These states can be experimentally observed as hot bands from the first excited torsional bands in the range 360 - 720 cm^{-1} . In the same range a hot band to the excited inversion state was discovered and since the intensities of three hot bands are similar, they must be analyzed jointly. The spectra were recorded with a resolution of 0.00125 cm^{-1} using Bruker IFS-120HR spectrometer at the University of Oulu. The accurate energy levels of the first excited torsional state, ν_{15} ,¹² were used as reference values for lower state combination differences in the assignments of the third and fourth torsional hot bands, $3\nu_{15} \leftarrow \nu_{15}$, $4\nu_{15} \leftarrow \nu_{15}$, as well as in the hot inversion-torsional band, $\nu_9 \leftarrow \nu_{15}$. On the basis of the predicted energy levels for the third and fourth excited torsional states, many transitions of the hot torsional bands: $3\nu_{15} \leftarrow \nu_{15}$, $4\nu_{15} \leftarrow \nu_{15}$, not assigned previously, have been identified. Many transitions belonging to the inversion-torsion hot band, $\nu_9 \leftarrow \nu_{15}$, have been assigned too. All the assignments were confirmed by the Lower State Combination Differences. Only experimental data of inversion-torsion hot band, $\nu_9 \leftarrow \nu_{15}$, could be fitted to a single state model based on the group theoretical formalism of Hougen and Ohashi³. The transitions belonging to the hot torsional bands ($3\nu_{15} \leftarrow \nu_{15}$, $4\nu_{15} \leftarrow \nu_{15}$) require another approach like a "hybrid" Hamiltonian model⁴⁵, thus the analysis is still in progress.

¹ I. Gulaczyk, M. Kreglewski, V-M. Horneman, *J. Mol. Spectrosc.* 342 (2017) 25-30.

² I. Gulaczyk, M. Kreglewski, *JQSRT* 252 (2020) 107097.

³ N. Ohashi and J.T. Hougen, *J. Mol. Spectrosc.* 121, (1987) 474.

⁴ I. Kleiner, J.T. Hougen, *J. Phys. Chem. A* 119 (2015) 10664.

⁵ I. Kleiner, J.T. Hougen, *J. Mol. Spectrosc.* 368 (2020) 111255.