

NEW LINE INTENSITIES FOR THE FAR INFRARED BANDS OF THE TRANS- AND CIS-CONFORMER OF NITROUS ACID (HONO), NEW DETERMINATION OF THE TRANS-CIS CONFORMER BARRIER AND ITS IMPACT ON THE ASTROPHYSICAL DETECTION OF NITROUS ACID IN PROTOSTELLAR CLOUDS

W. TCHANA BETNGA, F. KWABIA TCHANA, Université Paris Cité and Univ Paris Est Creteil, CNRS, LISA, F-75013 Paris, France; **A. PERRIN**, Laboratoire de Météorologie Dynamique/IPSL, UMR CNRS 8539, Ecole Polytechnique, Université Paris-Saclay, RD36, 91128 Palaiseau Cedex, France; **L. MANCERON**, Ligne AILES, Synchrotron SOLEIL, L'Orme des Merisiers, St-Aubin BP48, 91192 Gif-sur-Yvette Cedex, France; **J. VANDER AUWERA**, Spectroscopy, Quantum Chemistry and Atmospheric Remote Sensing (SQUARES), C.P. 160/09, Université Libre de Bruxelles, B-1050 Brussels, Belgium; **F. HINDLE**, Laboratoire de Physico-Chimie de l'Atmosphère, Université du Littoral Côte d'Opale, Dunkerque, France; **A. COUTENS**, Institut de Recherche en Astrophysique et Planétologie (IRAP), Université de Toulouse, UPS, CNRS, CNES, 9 av. du Colonel Roche, 31028 Toulouse Cedex 4, France

The first goal of this work is to improve the determination of the energy difference ($\Delta E_{Cis-Trans}$) between the ground vibrational state of the Cis- and Trans-HONO conformers of nitrous acid. For this, high resolution spectra were recorded in the 50-200 cm⁻¹ spectral region at three different temperatures, 240, 270, and 296 K. The relative line intensities for the B-type transitions of pure rotational bands of Trans-HONO and Cis-HONO achieved from our measurements were combined in least squares fit computations to those measured previously by Sironneau *et al.*¹. In this way, we could improve significantly the accuracy on the HONO conformer energy difference, with a value for $\Delta E_{Cis-Trans} = 95.8 \pm 9.2$ cm⁻¹ compared to $SIR \Delta E_{Cis-Trans} = 99 \pm 25$ cm⁻¹ in the previous study of Sironneau *et al.* The second goal is to generate a line list with absolute line intensities for the pure rotational bands in the far infrared region of Trans-HONO and Cis-HONO, with both A- and B-type transitions. This new line list proved to be more faithful for an improved detection of HONO in Astrophysical objects².

¹Sironneau V, Flaud JM, Orphal J, Kleiner I, Chelin P. Absolute line intensities of HONO and DONO in the far-infrared and re-determination of the Energy Difference between the trans- and cis-species of nitrous acid. *J Mol Spectrosc* 2010;**259**:100-104

²Coutens A, Ligerink NFW, Loison JC, Wakelam V, Calcutt H, Drozdovskaya MN, Jørgensen JK, Müller HSP, Van Dishoeck EF, Wampfler SF. The ALMA-PILS survey: First detection of nitrous acid (HONO) in the interstellar medium. *Astronomy & Astrophysics* 2019;**623**:L13