

## ROTATIONAL SPECTROSCOPY OF MOLECULAR IONS USING A NOVEL DOUBLE RESONANCE APPROACH

**O. ASVANY, P. C. SCHMID, W. SILVA, D. GUPTA, T. SALOMON, S.  
THORWIRTH, S. SCHLEMMER**, *I. Physikalisches Institut, Universität zu Köln,  
Zùlpicher Str. 77, 50937 Köln, Germany*

A novel method of rotational spectroscopy of trapped, cold molecular ions is presented. It is a double resonance scheme combining rotational excitation (in the mm-wave region) with rovibrational excitation (in the IR). The rovibrational excitation is detected by the recently developed leak-out spectroscopy (LOS) approach<sup>1,2</sup>. LOS is based on the vibrational-translational energy transfer process occurring during the collision of the IR excited ion with a noble gas atom, after which the ion may escape the trap ("leak out") so that it can be counted by a detector. LOS and the rotational scheme based on it are very sensitive and generally applicable to any cation or anion. This poster presents rotational spectra for  $\text{HC}_3\text{O}^+$ ,  $c\text{-C}_3\text{H}_2\text{D}^+$  and  $\text{H}_2\text{CCCH}^+$ .

---

<sup>1</sup>[doi:10.1021/acs.jpca.2c05767](https://doi.org/10.1021/acs.jpca.2c05767), P. C. Schmid, O. Asvany, T. Salomon, S. Thorwirth, and S. Schlemmer, *J. Phys. Chem. A*, **126**, 8111-8117, (2022).

<sup>2</sup>see also talk of P. C. Schmid in Minisymposium "Spectroscopy of ions in traps"