

## THE BENDING OF C<sub>3</sub>: EXPERIMENTALLY PROBING THE *l*-TYPE DOUBLING AND RESONANCE

**M.-A. MARTIN-DRUMEL**, **O. PIRALI**, **M. VERVLOET**, *Université Paris-Saclay, CNRS, Institut des Sciences Moléculaires d'Orsay, France*; **D. W. TOKARYK**, *Department of Physics, University of New Brunswick, Fredericton, NB, Canada*; **K. D. DONEY**, **H. LINNARTZ**, *Leiden Observatory, Laboratory for Astrophysics, Universiteit Leiden, Leiden, Netherlands*; **Q. ZHANG**, **Y. CHEN**, **D. ZHAO**, *Leiden Observatory, Laboratory for Astrophysics, Universiteit Leiden, Leiden, Netherlands*

C<sub>3</sub>, a pure carbon chain molecule that has been identified in different astronomical environments, is considered a good probe of kinetic temperatures through observation of transitions involving its low-lying bending mode ( $\nu_2$ ) in its ground electronic state. With the aim to investigate this bending mode with multiple quanta of excitation, we have undertaken high resolution optical and mid-infrared investigations on the species produced in discharge experiments.

We report here the most complete analysis of the  $\tilde{X}$  and  $\tilde{A}$  states of C<sub>3</sub> (literature and present study) using a single PGOPHER file. New experimental measurements result in 36 rovibronic  $\tilde{A}^1\Pi_u - \tilde{X}^1\Sigma_g^+$  bands (originating from  $\tilde{X}(0\nu_20)$ ,  $\nu_2 = 0 - 5$ , levels) recorded by laser induced fluorescence spectroscopy at the University of Science and Technology of China and the rovibrational  $\nu_3$  band (revealing hot bands involving up to 5 quanta of excitation in  $\nu_2$ ) recorded by Fourier-transform infrared spectroscopy using a global source on the AILES beamline of the SOLEIL synchrotron facility.

The combined fit allows for the accurate determination of the rotational parameters and absolute energy levels of C<sub>3</sub>, in particular for states involving the bending mode. The spectroscopic information derived from this work enables new interstellar searches for C<sub>3</sub>, not only in the infrared and optical regions investigated here but also notably in the  $\nu_2$  band region (around 63 cm<sup>-1</sup>) where vibrational satellites can now be accurately predicted. This makes C<sub>3</sub> a universal diagnostic tool to study very different astronomical environments, from dark and dense to translucent clouds.