FIRST ANALYSIS OF THE $0-\nu''$ PROGRESSION OF THE COMET-TAIL SYSTEM ${\bf A}^2\Pi_i-{\bf X}^2\Sigma^+$ IN THE $^{12}{\bf C}^{17}{\bf O}^+$ CATION.

I. PIOTROWSKA, W. SZAJNA, S. RYZNER, A. STASIK, A. PARA, P. KOLEK, R. KĘPA, R. HAKALLA, Materials Spectroscopy Laboratory, Institute of Physics, University of Rzeszów, Pigonia 1 Street, 35-310 Rzeszów, Poland; M.I. MALICKA, The Faculty of Mathematics and Applied Physics, Rzeszów University of Technology, Powstańców Warszawy 8 Street, 35-959 Rzeszów, Poland

The $0-\nu''$ progression of the Comet-Tail $A^2\Pi_i-X^2\Sigma^+$ system, where $\nu''=(0,1,2,3)$, was recorded for the first time in the $^{12}C^{17}O^+$ isotopologue. Using a high-resolution Fourier-transform spectrometer, more than 1500 emission lines were measured in the region of 12,880 - 20,500 cm $^{-1}$.

A deperturbation analysis was performed using the PGOPHER program.¹ As a result, deperturbed molecular parameters of the $A^2\Pi_i(\nu=0)$ and $X(\nu=0,1,2,3)$ levels and the $A^2\Pi_i(\nu=0) \sim X^2\Sigma^+(\nu=10)$ perturbation parameter of a complex (spin-orbit, spin-electronic and rotation-electronic) character were determined.

The values of the ro-vibronic terms of the $A^2\Pi_i(\nu=0)$ level were also calculated and the percentage $^2\Pi$ character of the $A^2\Pi_i(\nu=0)$ and $X^2\Sigma^+(\nu=10)$ levels was examined. This work is a continuation of the studies on the $A^2\Pi_i$ state in the $^{12}C^{17}O^+$ isotopologue made by our team.²

¹doi:10.1016/j.jqsrt.2016.04.010, C. M. Western, PGOPHER: a program for simulating rotational, vibrational and electronic spectra, *Journal of Quantitative Spectroscopy and Radiative Transfer*, **186**, 221-242, (2017).

 $^{^2}$ doi:10.1016/j.jqsrt.2022.108268, I. Piotrowska, R. Hakalla, W. Szajna, R. Kępa, First observation of the Comet-Tail $A^2\Pi_i - X^2\Sigma^+$ system in the $^{12}C^{17}O^+$ cation, *Journal of Quantitative Spectroscopy and Radiative Transfer*, **289**, 108268, (2022).