

ELECTRON SPIN-TORSION AND SPIN-ROTATION EFFECTS IN THE OPEN-SHELL CH₃CO RADICAL¹

L. H. COUDERT, O. PIRALI, R. CHAHBAZIAN, M.-A. MARTIN-DRUMEL,
Institut des Sciences Moléculaires d'Orsay, Université Paris-Saclay, Orsay, France;
L. ZOU, R. A. MOTIYENKO, L. MARGULÈS,
UMR 8523 - PhLAM - Physique des Lasers Atomes et Molécules, University of Lille, CNRS, F-59000 Lille, France

Although open-shell molecules displaying a fine interaction and non-rigid molecules displaying a large amplitude motion have been the subject of numerous spectroscopic investigations, fewer results are available about molecules displaying *both* a fine interaction and a large amplitude motion. A non-exhaustive list of such species include the open shell CH₂, CH₂OH, and Na₃ species displaying² respectively a large amplitude bending, torsion, and pseudorotation. The acetyl radical CH₃CO is a benchmark molecule for this subclass of molecules as it presents a low barrier internal rotation of its methyl group and a fine interaction. In the first spectroscopic investigation of the radical,³ a few low J , $K_a = 0$ transitions were measured in its microwave spectrum and a new theoretical model was developed to model its spectrum.

In this poster, new measurements carried out in the submillimeter wave spectrum of CH₃CO will be presented. More than 300 transitions have been assigned up to $N = 22$ and $K_a = 5$ in the ground torsional state for both A and E torsional species and for both F_1 and F_2 electron spin-rotation components. Based on a newly developed theoretical approach, a line frequency analysis has been carried out and line frequencies were reproduced with a unitless standard deviation of 1.2.

The analysis shows that two effects, ignored in the pioneering approach of Hirota *et al.*,³ are quite important. These are the electron spin-torsion coupling, described by a term in $p_\alpha S_z$, where p_α is the momentum conjugated to the torsional angle α , and a dependence of the electron spin-rotation coupling tensor on this torsional angle. In the poster the new measurements will be presented and the spectroscopic parameters will be reported. These include the usual torsion-rotation parameters A , B , C , D_{ab} , ρ , and V_3 , and the new fine interaction-torsion parameters.

¹Supported by the Programme National "Physique et Chimie du Milieu Interstellaire" (PCMI) of CNRS/INSU with INC/INP co-funded by CEA and CNES

²Ohashi, Tsuura, Hougen, Ernst, and Rakowsky, *J. Mol. Spec.* **184** (1997) 22; Coudert, *J. Chem. Phys.* **153** (2020) 144115; and Coudert, Chitarra, Spaniol, Loison, Martin-Drumel, and Pirali, *J. Chem. Phys.* **156** (2022) 244301

³Hirota, Mizoguchi, Ohsima, Katoh, Sumiyoshi, and Endo, *Mol. Phys.* **105** (2007) 455